

Overlap of Fractional Cloud for Radiation Calculations in GCMs: A Global Analysis using CloudSat and CALIPSO Data

H. W. Barker
Environment Canada

The need to consider cloud overlap structure depends on
the genre of the GCM:



Cloud Physics and Severe Weather Research Division
Environment Canada



Overlap of Fractional Cloud for Radiation Calculations in GCMs: A Global Analysis using CloudSat and CALIPSO Data

H. W. Barker
Environment Canada

The need to consider cloud overlap structure depends on the genre of the GCM:

- ***conventional GCMs:*** description of cloud overlap for unresolved cloud fields must be provided, via parametrization, to carry out, at least, radiative transfer calculations.



Cloud Physics and Severe Weather Research Division
Environment Canada



Overlap of Fractional Cloud for Radiation Calculations in GCMs: A Global Analysis using CloudSat and CALIPSO Data

H. W. Barker
Environment Canada

The need to consider cloud overlap structure depends on the genre of the GCM:

- **conventional GCMs:** *description of cloud overlap for unresolved cloud fields must be provided, via parametrization, to carry out, at least, radiative transfer calculations.*
- **MMF-GCMs (global CSRMs):** *cloud overlap ceases to be a parametrization issue and becomes a diagnostic variable*



*Cloud Physics and Severe Weather Research Division
Environment Canada*



Objectives

- make a global assessment of cloud overlap
 - using CloudSat-CALIPSO data
- estimate global-average radiative sensitivity for overlap
$$\frac{\partial F}{\partial \mathcal{L}_{cf}^*}$$
- assess feasibility of a very simple overlap *parametrization*

Data


CloudSat-CALIPSO cloud-mask: Jan and Aug 2007

- http://cloudsat.cira.colostate.edu/data_dist/OrderData.php
- 2B-GEOPROF and 2B-GEOPROF-LIDAR
- *CPR_Cloud_mask* ; *Radar_Reflectivity* ; *CloudFraction*
 - cross-sections: 50, 100, 200, 500, and 1000 km
 - what best represents a GCM column (Astin + Di Girolamo 1999)?
 - ~37,000 columns/orbit... many thousands of samples
 - total cloud fractions $\in [0.05, 0.99]$
 - CloudSat's radar reflectivity $\sim r^6$... precipitation-mask???

precipitation-mask


“cloud” in bins 3 and 4 above the surface?

remove up to $\max(R)$
when lidar indicates
no cloud



Methodology

Hogan and Illingworth (2000):



$$c_{k,l} = \alpha_{k,l} \underbrace{\max(c_k, c_l)}_{\text{maximum overlap}} + (1 - \alpha_{k,l}) \underbrace{(c_k + c_l - c_k c_l)}_{\text{random overlap}}$$

$$\alpha_{k,l} \equiv \exp \left[- \int_{z_k}^{z_l} \frac{dz}{\mathcal{L}_{cf}(z)} \right] \quad \Bigg| \quad \mathcal{L}_{cf}(z) \in [0, \infty) ; \alpha_{k,l} \in [0, 1)$$

$$c_{k,l} \in (\max(c_k, c_l), c_k + c_l - c_k c_l]$$

Effective Decorrelation Length



- assume \mathcal{L}_{cf} does not vary vertically... unique $C(\mathcal{L}_{cf})$ given cloud fraction profile
- total cloud fraction \hat{C}
from measurements
- using McICA's sub-grid
cloud generator, solve:

$$C(\mathcal{L}_{cf}^*) = \hat{C}$$


Effective Decorrelation Length

- assume \mathcal{L}_{cf} does not vary vertically... unique $C(\mathcal{L}_{cf})$ given cloud fraction profile
- total cloud fraction \hat{C} from measurements
- using McICA's sub-grid cloud generator, solve:

$$C(\mathcal{L}_{cf}^*) = \hat{C}$$




January 2007




almost independent of cross-section length except for very small lengths
(scale-independent parametrization?)

January 2007



- little difference between land and ocean
- linear increase with C for small C
- peak medians of 2 to 3 km near $C = 0.7$

500 km cross-sections




largest values in Polar areas during winter and N. Tropics during boreal summer

- sedimentation of crystals and convection?

500 km cross-sections

January



- MMF values are very large (4 km grid-spacing vs. ~1.5 km?)
- screen for precip: minor for CloudSat, major for MMF

On the use of \mathcal{L}_{cf}^* in GCMs

- is vertically-constant sufficient?
- how to set it?... not from total cloud fraction...
- how detailed need the parametrization be?
- is it something that changes with climate?...

$$F_{\text{ICA}} = \int_0^\infty p(\tau) F(\tau) d\tau$$

$$F_{\text{ICA}} = (1 - \hat{C}) F(0) + \hat{C} \int_0^\infty \hat{p}(\tau) F(\tau) d\tau$$


$$F_{\text{ICA}} = (1 - \hat{C}) F(0) + \sum_{m=1}^M \varepsilon_m \int_0^\infty \hat{p}_m(\tau) F(\tau) d\tau$$

total cloud fraction

fraction of clouds in layer m
with tops exposed to space

distribution of τ for clouds in layer
 m with tops exposed to space


January 2007



- achieving an accurate distribution of cloudtops exposed to space appears doable
(given *correct* cloud fraction profiles and effective decorrelation lengths)



500 km cross-sections

January 2007



- abundant (but significant?) random noise due to global setting
 - max-rand scheme has this *noise* too
- is it necessary to reduce variance?... cf. McICA

ECMWF



Radiative Sensitivity for Overlap

- to what extent is TOA radiation affected by overlap?...

Radiative Sensitivity for Overlap

- to what extent is TOA radiation affected by overlap?

$$F_{\text{ICA}} = (1 - \hat{C})F(0) + \hat{C} \int_0^\infty \hat{p}(\tau)F(\tau) d\tau = (1 - \hat{C})F(0) + \hat{C}F_{\text{cld}}$$

$$\frac{\partial F_{\text{ICA}}}{\partial \mathcal{L}_{cf}} = -F(0) \frac{\partial \hat{C}}{\partial \mathcal{L}_{cf}} + F_{\text{cld}} \frac{\partial \hat{C}}{\partial \mathcal{L}_{cf}} + \hat{C} \frac{\partial F_{\text{cld}}}{\partial \mathcal{L}_{cf}}$$

$$\frac{\partial F_{\text{ICA}}}{\partial \mathcal{L}_{cf}} \simeq CRE \frac{\partial \ln \hat{C}}{\partial \mathcal{L}_{cf}} + \hat{C} \left[\underbrace{\frac{\partial F_{\text{cld}}}{\partial \bar{\tau}} \frac{\partial \bar{\tau}}{\partial \mathcal{L}_{cf}}}_{< 0} + \underbrace{\frac{\partial F_{\text{cld}}}{\partial \sigma_\tau} \frac{\partial \sigma_\tau}{\partial \mathcal{L}_{cf}}}_{\sim O(0)} + \dots \right]$$

$$\begin{aligned} \left\langle \frac{\partial F_{\text{ICA}}}{\partial \mathcal{L}_{cf}} \right\rangle &\simeq \langle CRE \rangle \left\langle \frac{\partial \ln \hat{C}}{\partial \mathcal{L}_{cf}} \right\rangle - O(\lesssim 1) \\ &\simeq (-45 \text{ Wm}^{-2}) (-0.08 \text{ km}^{-1}) - O(\lesssim 1) \\ &\simeq 3 \text{ Wm}^{-2} \text{ km}^{-1} \end{aligned}$$

Radiative Sensitivity for Overlap

- to what extent is TOA radiation affected by overlap?

$$F_{\text{ICA}} = (1 - \hat{C})F(0) + \hat{C} \int_0^\infty \hat{p}(\tau)F(\tau) d\tau = (1 - \hat{C})F(0) + \hat{C}F_{\text{cld}}$$

$$\frac{\partial F_{\text{ICA}}}{\partial \mathcal{L}_{cf}} = -F(0) \frac{\partial \hat{C}}{\partial \mathcal{L}_{cf}} + F_{\text{cld}} \frac{\partial \hat{C}}{\partial \mathcal{L}_{cf}} + \hat{C} \frac{\partial F_{\text{cld}}}{\partial \mathcal{L}_{cf}}$$

$$\frac{\partial F_{\text{ICA}}}{\partial \mathcal{L}_{cf}} \simeq CRE \frac{\partial \ln \hat{C}}{\partial \mathcal{L}_{cf}} + \hat{C} \left[\underbrace{\frac{\partial F_{\text{cld}}}{\partial \bar{\tau}} \frac{\partial \bar{\tau}}{\partial \mathcal{L}_{cf}}}_{< 0} + \underbrace{\frac{\partial F_{\text{cld}}}{\partial \sigma_\tau} \frac{\partial \sigma_\tau}{\partial \mathcal{L}_{cf}}}_{\sim O(0)} + \dots \right]$$

$$\left\langle \frac{\partial F_{\text{ICA}}}{\partial \mathcal{L}_{cf}} \right\rangle \simeq \langle CRE \rangle \left\langle \frac{\partial \ln \hat{C}}{\partial \mathcal{L}_{cf}} \right\rangle - O(\lesssim 1)$$

$$\simeq (-45 \text{ Wm}^{-2}) (-0.08 \text{ km}^{-1}) - O(\lesssim 1)$$

$$\simeq 3 \text{ Wm}^{-2} \text{ km}^{-1} \quad (\text{LW is at least } \sim 3 \text{ times smaller})$$

Conclusions + Recommendations

- 2 months of overlap analyses... more is needed
- bring in additional data (e.g., ECMWF)
- is \mathcal{L}_{cf}^* sufficient?
 - can it be as simple as a few judicious settings based on local conditions?
- assess GCMs recognizing

$$F_{\text{ICA}} = (1 - \hat{C})F(0) + \sum_{m=1}^M \varepsilon_m \int_0^\infty \hat{p}_m(\tau) F(\tau) d\tau$$

- if these are correct, so too is overlap, and your radiation budget

January 2007

